



Europäisches Patentamt  
European Patent Office  
Office européen des brevets



(11) **EP 0 953 339 A1**

(12) **EUROPEAN PATENT APPLICATION**

(43) Date of publication:  
03.11.1999 Bulletin 1999/44

(51) Int. Cl.<sup>6</sup>: **A61K 7/48, C07F 9/10**

(21) Application number: **99105069.1**

(22) Date of filing: **23.03.1999**

(84) Designated Contracting States:  
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU  
MC NL PT SE**  
Designated Extension States:  
**AL LT LV MK RO SI**

(72) Inventors:  
- **Takekoshi, Yoichiro**  
Tokyo (JP)  
- **Matsuura, Ichiro**  
Tokyo (JP)

(30) Priority: **24.03.1998 JP 7562498**

(74) Representative:  
**VOSSIUS & PARTNER**  
Siebertstrasse 4  
81675 München (DE)

(71) Applicant:  
**KYOWA HAKKO KOGYO CO., LTD.**  
Chiyoda-ku, Tokyo 100-8185 (JP)

(54) **Cosmetic composition**

(57) A composition for cosmetics use which comprises a lysophospholipid mixture, wherein 30 mol% or more of fatty acids bonded to said lysophospholipid mixture are monoenoic fatty acids; and a cosmetic composition which comprises a lysophospholipid mixture and a cosmetically acceptable carrier, wherein 30 mol% or more of fatty acids bonded to said lysophospholipid mixture are monoenoic fatty acids. The cosmetic composition having such a lysophospholipid mixture has high storage stability and is excellent in feeling upon use.

BEST AVAILABLE COPY

EP 0 953 339 A1

## Description

[0001] This invention relates to lysophospholipids for use in cosmetics having high storage stability and excellent in feeling upon use as well as to a cosmetic composition comprising the same.

[0002] Phospholipid is known as a surface active agent or emulsifying agent which is mild to the skin. Lysophospholipid obtained by converting the phospholipid into its lyso form by enzyme treatment or the like means is also known as a surface active agent or emulsifying agent which is tender to the skin (JP-A-61-112007, JP-A-63-41411; the term "JP-A" as used herein means a "Japanese published unexamined patent application"). Examples of the known lysophospholipid include those which are produced from phospholipid of animal origin such as the egg yolk, bovine brain or swine brain or from phospholipid of plant origin such as soybean, corn, rapeseed or the like plant seed, and they are used in cosmetics, quasi drugs, toiletries and the like, as surface active agents, solubilizing agents, emulsifying agents, gelling agents, moisture-keeping agents or liposome-making agents.

[0003] Since lysophospholipid, when compounded in cosmetics, can provide the products with a soft touch, it is an effective material for improving the touch of products during and after their use.

[0004] However, since there are problems from the viewpoint of stability (such as odor and discoloration), there is a limitation, in terms of formulation designing such as blending amount and masking of an oxidized soybean-like odor. Although a hydrogenated lysolactin is available (in which unsaturated fatty acids of lysophospholipid are hydrogenated) in order to improve stability, the material provides poor feel because of less adaptability and moist sensation on the skin.

[0005] The object of the present invention is to provide lysophospholipids for use in cosmetics having both high storage stability and excellent feeling upon use, a cosmetic composition comprising the same and methods using the lysophospholipid.

[0006] This object has been achieved by the surprising finding that cosmetics having excellent stability and feeling upon use can be obtained by preparing cosmetic compositions with lysophospholipid obtained by converting high oleic phospholipid into its lyso form.

[0007] The present invention relates to the following (1) to (14).

(1) A composition comprising lysophospholipids, wherein at least 30 mol% of the fatty acids bonded to said lysophospholipids are monoenoic fatty acids.

(2) The composition according to (1), wherein at least 50 mol% of the fatty acids bonded to said lysophospholipids are monoenoic fatty acids.

(3) The composition according to (2), wherein at least 70 mol% of the fatty acids bonded to said lysophospholipids are monoenoic fatty acids.

(4) The composition according to any one of (1) to (3), wherein at most 20 mol% of the fatty acids bonded to said lysophospholipids are polyenoic fatty acids.

(5) The composition according to (4), wherein at most 15 mol% of the fatty acids bonded to said lysophospholipids are polyenoic fatty acids.

(6) The composition according to (5), wherein at most 10 mol% of the fatty acids bonded to said lysophospholipids are polyenoic fatty acids.

(7) The composition according to any one of (1) to (6), wherein said monoenoic fatty acids have 4-34 carbon atoms.

(8) The composition according to (7), wherein said monoenoic fatty acids have 14-24 carbon atoms.

(9) The composition according to any one of (1) to (8), wherein saturated fatty acids bonded to said lysophospholipids have 1-34 carbon atoms.

(10) The composition according to (9), wherein saturated fatty acids bonded to said lysophospholipids have 6-24 carbon atoms.

(11) A cosmetic composition, which comprises the composition according to any one of (1) to (10) and a cosmetically acceptable carrier.

(12) A method of improving storage stability in a cosmetic composition, which comprises the step of blending lysophospholipids according to any one of (1) to (10) into a cosmetic composition comprising cosmetically acceptable carrier, wherein said lysophospholipids are present in said cosmetic composition in a total amount of 0.01 to 99.99% by weight.

(13) A method of improving feeling upon use of a cosmetic composition, which comprises the step of blending lysophospholipids according to any one of (1) to (10) into a cosmetic composition comprising cosmetically acceptable carrier, wherein said lysophospholipids are present in said cosmetic composition in a total amount of 0.01 to 99.99% by weight.

(14) A method of providing suitable feeling upon use of a cosmetic composition, which comprises the step of applying lysophospholipids according to any one of (1) to (10) to the body, wherein said lysophospholipids are present

in said cosmetic composition in a total amount of 0.01 to 99.99% by weight.

[0008] The lysophospholipid of the present invention is glycerol-based. In this event, the lysophospholipid may be either 1-acyl-2-lysophospholipid or 2-acyl-1-lysophospholipid or a mixture thereof.

[0009] Kinds of the phospholipid are not particularly limited, and examples thereof include phosphatidylcholine, phosphatidylethanolamine, phosphatidylinositol, phosphatidic acid, phosphatidylglycerol, N-acylphosphatidylethanolamine, phosphatidylserine and the like which may be used alone or as a mixture thereof.

[0010] Preferred examples of the phospholipid to be used in the present invention include phosphatidylcholine, phosphatidylethanolamine, phosphatidylinositol and phosphatidic acid.

[0011] Regarding the fatty acids bonded to the lysophospholipid mixture, those in which 30 mol% or more of the total fatty acids are monoenoic fatty acids are desirable, 50 mol% or more are monoenoic fatty acids are more desirable and 70 mol% or more are monoenoic fatty acids are most desirable.

[0012] The remaining fatty acids are saturated fatty acids and polyenoic fatty acids. Regarding the fatty acids bonded to the lysophospholipid mixture to be used in the present invention, those in which 30 mol% or more of the total fatty acids are monoenoic fatty acids and the polyenoic fatty acid content is 20 mol% or less are desirable, 30 mol% or more of the total fatty acids are monoenoic fatty acids and the polyenoic fatty acid content is 15 mol% or less are more desirable and 30 mol% or more of the total fatty acids are monoenoic fatty acids and the polyenoic fatty acid content is 10 mol% or less are most desirable.

[0013] The monoenoic fatty acid is a fatty acid which has one double bond in the alkyl chain moiety of a saturated fatty acid. Though the number of its carbon atoms is not particularly limited, straight or branched monoenoic fatty acids having 4 to 34 carbon atoms can be exemplified, and monoenoic fatty acids having 14 to 24 carbon atoms are desirable. Illustrative examples of the monoenoic fatty acid include butenoic acid, pentenoic acid, hexenoic acid, heptenoic acid, octenoic acid, nonenoic acid, decenoic acid, undecenoic acid, dodecenoic acid, tridecenoic acid, tetradecenoic acid, pentadecenoic acid, hexadecenoic acid, heptadecenoic acid, octadecenoic acid, nonadecenoic acid, icosanoic acid, heneicosanoic acid, docosanoic acid, tricosanoic acid, tetracosanoic acid, pentacosanoic acid, hexacosanoic acid, heptacosanoic acid, octacosanoic acid, nonacosanoic acid, triacontanoic acid, hentriacontanoic acid, dotriacontanoic acid, tritriacontanoic acid, tetratriacontanoic acid and the like.

[0014] The decenoic acid includes obtusilic acid and the like, the dodecenoic acid includes linderic acid and the like, the tetradecenoic acid includes tsuzuic acid and the like, the hexadecenoic acid includes palmitoleic acid and the like and the octadecenoic acid includes petroselinic acid, oleic acid, elaidic acid, vaccenic acid and the like.

[0015] Any fatty acid may be used as the monoenoic fatty acid, with the proviso that it has one double bond, and it may be used as a single monoenoic fatty acid or as a mixture of monoenoic fatty acids having different carbon numbers or, even in the case of the same number of carbon atoms, monoenoic fatty acids having different position of the double bond. Oleic acid is desirable as the monoenoic fatty acid.

[0016] Regarding the fatty acid bonded to the lysophospholipid mixture to be used in the present invention, any fatty acid may be bonded, with the proviso that it satisfies the aforementioned conditions.

[0017] With respect to the saturated fatty acid, the number of carbon atoms is not particularly limited, and straight or branched saturated fatty acids having 1 to 34 carbon atoms can be exemplified, of which saturated fatty acids having 6 to 24 carbon atoms are desirable. Illustrative examples of the saturated fatty acid include formic acid, acetic acid, propionic acid, butanoic acid (butyric acid), pentanoic acid, hexanoic acid (caproic acid), heptanoic acid, octanoic acid (caprylic acid), nonanoic acid, decanoic acid (capric acid), undecanoic acid, dodecanoic acid (lauric acid), tridecanoic acid, tetradecanoic acid (myristic acid), pentadecanoic acid, hexadecanoic acid (palmitic acid), heptadecanoic acid (margaric acid), octadecanoic acid (stearic acid), nonadecanoic acid, icosanoic acid (arachidic acid), heneicosanoic acid, docosanoic acid (behenic acid), tricosanoic acid, tetracosanoic acid (lignoceric acid), pentacosanoic acid, hexacosanoic acid (cerotic acid), heptacosanoic acid, octacosanoic acid (montanolic acid), nonacosanoic acid, triacontanoic acid (melleic acid), hentriacontanoic acid, dotriacontanoic acid, tritriacontanoic acid, tetratriacontanoic acid and the like.

[0018] The polyenoic fatty acid is a fatty acid which has two or more double bonds in the alkyl chain moiety of a saturated fatty acid, and its examples include a dienoic fatty acid which has two double bonds in the alkyl chain moiety of a saturated fatty acid and a trienoic fatty acid which has three double bonds in the alkyl chain moiety of a saturated fatty acid. The position of double bonds in the alkyl moiety of the polyenoic fatty acid is not particularly limited.

[0019] Illustrative examples of the dienoic fatty acid include straight or branched fatty acid having 6 to 34 carbon atoms such as hexadienoic acid, heptadienoic acid, octadienoic acid, nonadienoic acid, decadienoic acid, undecadienoic acid, dodecadienoic acid, tridecadienoic acid, tetradecadienoic acid, pentadecadienoic acid, hexadecadienoic acid, heptadecadienoic acid, octadecadienoic acid, nonadecadienoic acid, icosadienoic acid, heneicosadienoic acid, docosadienoic acid, tricosadienoic acid, tetracosadienoic acid, pentacosadienoic acid, hexacosadienoic acid, heptacosadienoic acid, octacosadienoic acid, nonacosadienoic acid, triacontadienoic acid, hentriacontadienoic acid, dotriacontadienoic acid, tritriacontadienoic acid, tetratriacontadienoic acid and the like. The octadecadienoic acid includes

linoleic acid.

[0020] Illustrative examples of the trienoic fatty acid include straight or branched fatty acids having 8 to 34 carbon atoms such as octatrienoic acid, nonatrienoic acid, decatrienoic acid, undecatrienoic acid, dodecatrienoic acid, tridecatrienoic acid, tetradecatrienoic acid, pentadecatrienoic acid, hexadecatrienoic acid, heptadecatrienoic acid, octadecatrienoic acid, nonadecatrienoic acid, icosatrienoic acid, heneicosatrienoic acid, docosatrienoic acid, tricosatrienoic acid, tetracosatrienoic acid, pentacosatrienoic acid, hexacosatrienoic acid, heptacosatrienoic acid, octacosatrienoic acid, nonacosatrienoic acid, triacontatrienoic acid, hentriacontatrienoic acid, dotriacontatrienoic acid, tritriacontatrienoic acid, tetratriacontatrienoic acid and the like. The octadecatrienoic acid includes, for example,  $\gamma$ -linolenic acid and  $\alpha$ -linolenic acid.

[0021] Fatty acids having four or more double bonds, such as arachidonic acid, eicosapentaenoic acid, docosahexaenoic acid and the like, can also be exemplified as the polyenoic acid.

[0022] The fatty acid to be bonded to the lysophospholipid may be substituted by one to four hydroxy groups or one or two oxy groups.

[0023] The lysophospholipid to be used in the present invention can be produced by synthesis, but it is desirable to produce it by enzymatically hydrolyzing an acyl group at a specific site of natural phospholipid or synthesized phospholipid with phospholipase A<sub>1</sub> or phospholipase A<sub>2</sub>. As the phospholipid, natural phospholipid is desirable because it can be obtained with a low cost. The phospholipid may be used by modifying it in advance by its treatment with phospholipase D or the like enzyme, or by fractionating its specific components using a solvent or the like.

[0024] It is desirable that, after the enzyme treatment, the lysophospholipid to be used in the present invention is purified by further treating it with acetone, methanol, ethanol, propanol, isopropanol and the like organic solvents.

[0025] Animal or plant phospholipid can be exemplified as the natural phospholipid. Examples of the plant phospholipid include those which are obtained from high oleic safflower, high oleic corn, lauric acid rich rapeseed and the like. As other sources of phospholipid, it is possible to use a plant phospholipid whose fatty acid composition is modified to the aforementioned composition for example by genetic recombination techniques. Egg yolk phospholipid can be exemplified as the animal phospholipid. The lysophospholipid to be used in the present invention is obtained preferably by enzymatically treating a natural phospholipid mixture containing 80 mol% or more of monoenoic fatty acid with the aforementioned phospholipase, more preferably by enzymatically treating a natural phospholipid mixture containing 30 mol% or more of monoenoic fatty acid and 20 mol% or less of polyenoic fatty acid with the aforementioned phospholipase.

[0026] Isolation of phospholipid from an animal or plant material can be made by any method. For example, the gum formed by adding water to crude oil at the degumming step during the process of plant oil production can be used.

[0027] Examples of the cosmetics of the present invention include toilet soaps, cleansing articles, shampoos, rinses, hairdyes, hair cosmetics, creams, milky lotions, face lotion, face oils, sunburn/suntan oils, face powders/powders, pecks, nail creams, cheek rouges, eyebrow pencils, eye creams, eye shadows, mascara articles, lipsticks, lip creams, dental creams, bathing cosmetics and the like.

[0028] Though not particularly limited, examples of the body part to which the cosmetics of the present invention are applied include head hair, body hair, skin, face, lips, mouth and the like.

[0029] When the cosmetics of the present invention are applied to the body, excellent feeling upon use such as softness, spreading touch or conformability is obtained during their use, and excellent feeling upon use such as moistness or refreshment is obtained after their use.

[0030] The cosmetic of the present invention comprises lysophospholipid and a carrier acceptable for cosmetics. The lysophospholipid content is preferably from 0.001 to 99.99% by weight, more preferably from 0.01 to 50% by weight, most preferably from 0.1 to 5% by weight.

[0031] Examples of the carrier include those materials which are generally used in cosmetics and pharmaceutical preparations, such as oils and fats, hydrocarbons, waxes, fatty acids, synthetic esters, alcohols, surface active agents, thickeners, moisture-keeping agents, antiseptics, antioxidants, pH adjusting agents, perfumes, pigments, drugs, purified water and the like, and these carriers may be used alone or as a mixture of two or more.

[0032] These carriers, alone or as a mixture of two or more, may be used in an amount of preferably from 0.001 to 99.99% by weight, more preferably from 50 to 99.99% by weight, most preferably from 95 to 99.5% by weight.

[0033] Examples of the oils and fats include jojoba oil, castor oil, olive oil, soybean oil, palm oil, cacao butter, camellia oil, coconut oil, Japan wax, grape seed oil, avocado oil, yolk oil, mink whale oil, turtle oil and the like.

[0034] Examples of the hydrocarbons include liquid paraffin, vaseline, microcrystalline wax, ceresin wax, paraffin wax, squalane,  $\alpha$ -olefin oligomer and the like.

[0035] Examples of the waxes include beeswax, lanolin, carnauba wax, candelilla wax, whale wax and the like.

[0036] Examples of the fatty acids include lauric acid, myristic acid, palmitic acid, stearic acid, oleic acid, isostearic acid, behenic acid and the like.

[0037] Examples of the synthetic esters include cetyl palmitate, isopropyl myristate, isopropyl palmitate, butyl oleate, octyldodecyl oleate, cholesterol oleate, myristyl myristate, octyldodecyl myristate, propylene glycol monooleate, myristyl

lactate, isoestearyl malate, glycerol monoostearate, distearyl dimethyl ammonium chloride and the like.

[0038] It is desirable that each of oils and fats, hydrocarbons, waxes, fatty acids and synthetic esters is blended to a ratio of from 0 to 70% by weight.

[0039] Examples of the alcohols include ethanol, 1,3-butylene glycol, propylene glycol, lauryl alcohol, cetanol, hexyl decanol, octyl dodecanol, stearyl alcohol, oleyl alcohol and the like. These alcohols are blended at a ratio of from 0 to 60% by weight.

[0040] Examples of the surface active agents include decaglycerolmonomyristate or the like lipophilic glycerol fatty acid ester, self emulsification-type glycerol fatty acid ester, polyglycerol fatty acid ester, polyoxyethylene glycerol fatty acid ester, polyoxyethylene hydrogenated castor oil, polyoxyethylene sorbitol fatty acid ester, polyoxyethylene (20) cetyl ether or the like polyoxyethylene alkyl ether, sodium polyoxyethylene fatty acid ester alkylsulfate, sodium polyoxyethylene alkylsulfate, sodium alkylbenzenesulfonate, polyoxyethyleneglyceryl pyroglutamate isostearate, dialkyl sulfosuccinate, cetyl pyridinium bromide, n-octadecyltrimethylammonium chloride, monoalkylphosphoric acid, N-acylglutamic acid, polyoxyethylene reduced lanolin and the like. It is desirable to blend these surface active agents at a ratio of from 0 to 40% by weight.

[0041] Examples of the thickeners include sodium alginate, xanthan gum, aluminum silicate, carboxyvinyl polymer, polyvinyl alcohol, methyl polysiloxane, dextran, carboxymethylcellulose, carrageenan, hydroxypropylmethylcellulose, soluble starch, cationated cellulose, tragacanth gum, quince seed extract and the like. It is desirable to blend these thickeners at a ratio of from 0 to 0.5% by weight.

[0042] Examples of the moisture-keeping agents include glycerol, propylene glycol, 1,3-butylene glycol, pyrrolidone-carboxylic acid, lactic acid, hyaluronic acid and the like. It is desirable to blend these moisture-keeping agents at a ratio of from 0 to 60% by weight.

[0043] Examples of the antiseptics include benzoic acid, salicylic acid or dehydroacetic acid or salts thereof, paraoxybenzoic acid ester such as methylparaben or propylparaben, triclosan halocarbon, benzalkonium chloride, hinokitol, resorcinol and the like. It is desirable to blend these antiseptics at a ratio of from 0 to 1% by weight.

[0044] Examples of the antioxidants include dibutylhydroxytoluene, butylhydroxyanisole, propyl gallate, ascorbic acid and the like. It is desirable to blend these antioxidants at a ratio of from 0 to 5% by weight.

[0045] Examples of the pH adjusting agents include sodium hydroxide, triethanolamine, citric acid, sodium citrate, boric acid, borax, potassium hydrogenphosphate and the like. It is desirable to blend these pH adjusting agents at a ratio of from 0 to 5% by weight.

[0046] Regarding the perfumes, any perfume can be used with the proviso that it is generally used in cosmetics.

[0047] Examples of the pigments include iron oxide, titanium dioxide, zinc oxide, kaolin, talc and the like. It is desirable to blend these pigments at a ratio of from 0 to 90% by weight.

[0048] Examples of the drugs include wheat germ oil, vitamin E, vitamin A, vitamin B<sub>2</sub>, ascorbic acid 2-phosphate magnesium salt, D-panthothenyl alcohol, dipotassium glycyrrhetinate, glutathione and the like. It is desirable to blend these drugs at a ratio of from 0 to 5% by weight.

#### TEST EXAMPLE 1

[0049] An organoleptic test was carried out using the face lotions produced in Example 1 and Comparative Example 1.

[0050] Softness and spreading/conformability during their use and moistness after use were evaluated as organoleptic characteristics by 10 cosmetics panels. The evaluation was carried out by a five point method, and the results were expressed as average values. The results are shown in Table 1.

Table 1

Items evaluated			Example 1	Comparative Example 1
Organoleptic characteristics	During use	Softness	5	2
		Spreading/ conformability	5	2
	After use	Moistness	5	3

[0051] The hydrogenated soybean lysophospholipid showed inferior softness and spreading/conformability during use and moistness after use when compared with the soybean lysophospholipid which, therefore, was excellent in feeling upon use.

**TEST EXAMPLE 2**

[0052] Organoleptic characteristics and emulsifying stability were evaluated using the face lotions produced in Example 1 and Comparative Example 2.

[0053] Softness and spreading touch during their use and refreshing feeling after use were evaluated as organoleptic characteristics by 10 cosmetic panels. The evaluation was carried out by a five point method, and the results were expressed as average values. The odor was judged as "good" when it did not change, "slightly good" when slightly changed or "bad" when markedly changed.

[0054] The storage stability test was carried out by an organoleptic inspection and an instrumental analysis after 1 to 2 weeks of storage of each sample at 5°C or 40°C or under UV irradiation (light source: BBL; FL20S-BL-B manufactured by Matsushita Electric Works, 20 cm in irradiation distance, room temperature) or sunlight irradiation.

[0055] Characteristics of the appearance were observed, and the results were judged as "good" when no changes were found, "slightly good" when slight changes were found or "bad" when significant changes were found.

[0056] Changes in color tone were evaluated using samples stored under UV irradiation or at 40°C, by measuring color differences from samples stored at room temperature. As the colorimeter, SE 2000 (Japan Denshoku Kogyo) was used.

[0057] The results are shown in Table 2.

Table 2

Items tested			Example 1	Comparative Example 2
Emulsification stability				
	5°C		good	good
	Room temperature		good	slightly good
	40°C		good	bad
	UV irradiation		slightly good	bad
Organoleptic characteristics				
	Odor	5°C	good	slightly good
		40°C	slightly good	bad
		UV irradiation	good	bad
	During use	Softness	5	3
		Spreading touch	5	3
	After use	Refreshing feeling	5	1
Color tone (ΔE)		Sunlight irradiation	-	-
		UV irradiation	0.48	2.24

[0058] Since the soybean lysophospholipid was sensitive to high temperature and light, it showed a sticky feeling as the sense of touch, and its appearance was considerably discolored by the UV irradiation. Regarding the high oleic safflower lysophospholipid, it was stable against temperature and light, so that its organoleptic inspection resulted in excellent softness, spreading touch and refreshing feeling. Changes in color tone were not found.

**TEST EXAMPLE 3**

[0059] Organoleptic characteristics and emulsifying stability were evaluated using the creams produced in Example 2 and Comparative Example 3.

[0060] The storage stability test was carried out by an organoleptic inspection and an instrumental analysis after 1 to 2 weeks of storage of each sample at 5°C or 40°C or under UV irradiation (light source: BBL; FL20S-BL-B manufactured by Matsushita Electric Works, 20 cm in irradiation distance, room temperature) or sunlight irradiation.

[0061] Characteristics of the appearance were observed, and the results were judged as "good" when no changes were found, "slightly good" when slight changes were found or "bad" when significant changes were found.

[0062] Changes in color tone were evaluated using samples stored under UV irradiation or at 40°C, by measuring

EP 0 953 339 A1

color differences from samples stored at room temperature. As the colorimeter, SE 2000 (Japan Denshoku Kogyo) was used.

[0063] The results are shown in Table 3.

Table 3

Items tested		Example 2	Comparative Example 3
Emulsification stability			
	5°C	good	good
	Room temperature	good	slightly good
	40°C	good	bad
	UV irradiation	slightly good	bad
Organoleptic characteristics			
	Odor	5°C	3
		40°C	4
		UV irradiation	5
	During use	Softness	5
		Spreading touch	5
	After use	Refreshing feeling	5
Color tone ( $\Delta E$ )			
		Sunlight irradiation	1.22
		UV irradiation	0.27
			5.97
			1.77

[0064] With regard to the amount of lipid peroxide, the peroxide value was obtained by the chloroform method [*Oil Chemistry*, 36, 276 - 278 (1987)], and the amount of thiobarbituric acid (TBA) reaction product was obtained by the TBA method [*Journal of the Japanese Society of Scientific Fisheries*, 45, 499 (1979)].

[0065] The results are shown in Table 4.

Table 4

Items tested		Example 2	Comparative Example 3
Peroxide value (chloroform method) (meq/kg)	Room temp.	6.5	6.1
	40°C	8.8	8.3
	UV irradiation	4.8	17.9
TBA reaction product (nmol/g)	Room temp.	1	3
	40°C	2	4
	UV irradiation	2	10

[0066] Since the soybean lysophospholipid was sensitive to high temperature and light, a sticky feeling remained as the sense of touch, and its appearance was considerably discolored by the sunlight and UV irradiation.

[0067] Regarding the high oleic safflower lysophospholipid, it was stable against temperature and light, so that its organoleptic inspection resulted in excellent spreading touch and conformability. Though color tone was changed by sunlight irradiation, it was not a coloration but a color difference changed from yellow to white. Changes in color tone were not found by UV irradiation.

[0068] Being sensitive to light, the soybean lysophospholipid showed accelerated oxidation in terms of both peroxide value and amount of TBA reaction product, which coincided with the results of organoleptic test on odor and color dif-

ference in coloration. The cosmetic formulated using high oleic lysophospholipid was stable against temperature and light and therefore was superior to the conventional soybean lysophospholipid.

[0069] Examples of the present invention are given below by way of illustration and not by way of limitation.

#### EXAMPLE 1

[0070] The contents of the following component (A) were mixed and dissolved and warmed up to 60°C, and the contents of the following component (B) mixed and dissolved at 55°C in advance were added to the component (A) and uniformly dissolved therein to produce a face lotion having the following composition. In this case, the high oleic safflower lysophospholipid produced in Example 3 was used. The amount of each compound was shown by % by weight.

(A)	Methylparaben (mfd. by Midori Kagaku)	0.1
	1,3-Butylene glycol (1,3-Butylene Glycol P, mfd. by Kyowa Yuka)	3.0
	Glycerol (conc. glycerol, mfd. by Sakamoto Pharmaceutical)	1.8
	Dipotassium glycyrrhetinate (Glycyron K <sub>2</sub> , mfd. by Tokiwa Phytochemistry Lab.)	0.1
	1% Sodium hyaluronate (mfd. by Kowa Hako Kogyo)	2.0
	Purified water	91.0
(B)	High oleic safflower lysophospholipid	0.2
	Propylene glycol (mfd. by Dow Chemical)	1.8

#### COMPARATIVE EXAMPLE 1

[0071] A face lotion was produced in the same manner as described in Example 1, except that hydrogenated soybean lysophospholipid (Hydrogenated Soybean lysophospholipid, mfd. by The Nisshin Oil Mills) was used instead of the high oleic safflower lysophospholipid.

#### COMPARATIVE EXAMPLE 2

[0072] A cream was produced in the same manner as described in Example 2, except that soybean phospholipid (mfd. by Kyowa Hako Kogyo) was used instead of the high oleic safflower lysophospholipid.

#### EXAMPLE 2

[0073] The contents of the following component (A) were mixed and dissolved at 85°C, the contents of the following component (B) mixed and dissolved in advance at 85°C were added to the component (A) and uniformly stirred, and then the following component (C) was added to the resulting mixture and uniformly mixed and cooled to produce a cream having the following composition. In this case, the high oleic safflower lysophospholipid produced in Example 3 was used. The amount of each compound was shown by % by weight.



5 10 15	(A)	Decaglyceryl monomyristate (NIKKOL Decaglyn, mfd. by Nikko Chemicals)	1.2
		POE (20) cetyl ether (EMALEX 1620, mfd. by Nippon Emulsion)	1.8
		$\alpha$ -Olefin oligomer (NIKKOL Syntheran 4, mfd. by Nikko Chemicals)	8.0
		Squalane (mfd. by Kishimoto Liver Oil)	4.0
		Cetyl palmitate (NIKKOL N-SP, mfd. by Nikko Chemicals)	3.0
		Cetanol (Conol 30 FC, mfd. by New Japan Chemical)	4.4
		Propylparaben (mfd. by Midori Kagaku)	0.03
		High oleic safflower lysophospholipid	2.0
20	(B)	Methylparaben (mfd. by Midori Kagaku)	0.1
		1,3-Butylene glycol (1,3-Butylene Glycol P, mfd. by Kyowa Yuka)	6.0
		Glycerol (conc. glycerol, mfd. by Sakamoto Pharmaceutical)	3.0
		Purified water	56.47
	(C)	0.5% Carboxyvinyl polymer solution (Carbopol 980, mfd. by B.F. Goodrich)	10.0

**COMPARATIVE EXAMPLE 3**

[0074] A cream was produced in the same manner as described in Example 2, except that soybean phospholipid (mfd. by Kyowa Hakko Kogyo) was used instead of the high oleic safflower lysophospholipid.

**EXAMPLE 3**

[0075] A 4 ml portion of 5N sodium hydroxide aqueous solution was added to 400 g of hydrated phospholipid having a moisture content of 55% (mfd. by True Lecithin Industry) obtained from the degumming step of high oleic safflower oil and the mixture was warmed up to 50°C, and then 24 hours of lyso-conversion reaction was carried out at 50°C by adding 0.12 ml of Lecithase (mfd. by NOVO) as a hydrolase to the stirring mixture.

[0076] Using a rotary evaporator, the reaction solution was concentrated and dehydrated with heating under a reduced pressure. Thereafter, diatomaceous earth was added to the resulting residue and the precipitate was removed using a pressure filtration apparatus, thereby obtaining an enzyme-treated phospholipid preparation.

[0077] Said treated preparation was washed with 10 volumes of acetone and dried under a reduced pressure to obtain an enzyme-treated purified phospholipid preparation (80 mol% in lyso-conversion ratio) in the form of powder.

**REFERENCE EXAMPLE 1**

[0078] Results of the fatty acid analysis of the high oleic safflower lysophospholipid produced in Example 3, the hydrogenated soybean lysophospholipid used in Comparative Example 1 and the soybean lysophospholipid used in Comparative Examples 2 and 3 are shown in Table 5. In this connection, the fatty acid composition was analyzed by firstly carrying out methyl esterification of each sample by the boron trifluoride-methanol method of the Standard Test Methods for Fat Analysis and then testing the resulting sample based on a flame ionization type constant temperature gas chromatography. Data in the table are shown by mol% of fatty acids.

Table 5

Fatty acids	High oleic safflower lyso-phospholipid	Soybean lysophospho-lipid	Hydrogenated soybean lysophospholipid
Palmitic acid	10.2	28.7	17
Stearic acid	1.2	6.2	57
Oleic acid	77.7	13.9	25
Linoleic acid	10.3	49.1	1
Linolenic acid	0.6	4.1	N.D.

N.D.: not detectable

[0079] As shown above, stable cosmetics having excellent feeling upon use can be provided by the present invention.

[0080] While the invention has been described in detail and with reference to specific embodiments thereof, it will be apparent to one skilled in the art that various changes and modifications can be made therein without departing from the spirit and scope thereof.

[0081] This application is based on Japanese patent application No. Hei. 10-75824, filed on March 24, 1998, and incorporated herein by reference.

#### Claims

1. A composition comprising lysophospholipids, wherein at least 80 mol% of the fatty acids bonded to said lysophospholipids are monoenoic fatty acids.
2. The composition according to claim 1, wherein at least 50 mol% of the fatty acids bonded to said lysophospholipids are monoenoic fatty acids.
3. The composition according to claim 2, wherein at least 70 mol% of the fatty acids bonded to said lysophospholipids are monoenoic fatty acids.
4. The composition according to any one of claims 1 to 3, wherein at most 20 mol% of the fatty acids bonded to said lysophospholipids are polyenoic fatty acids.
5. The composition according to claim 4, wherein at most 15 mol% of the fatty acids bonded to said lysophospholipids are polyenoic fatty acids.
6. The composition according to claim 5, wherein at most 10 mol% of the fatty acids bonded to said lysophospholipids are polyenoic fatty acids.
7. The composition according to any one of claims 1 to 6, wherein said monoenoic fatty acids have 4-34 carbon atoms.
8. The composition according to claim 7, wherein said monoenoic fatty acids have 14-24 carbon atoms.
9. The composition according to any one of claims 1 to 8, wherein saturated fatty acids bonded to said lysophospholipids have 1-34 carbon atoms.
10. The composition according to claim 9, wherein saturated fatty acids bonded to said lysophospholipids have 6-24 carbon atoms.
11. A cosmetic composition, which comprises a composition comprising lysophospholipids as defined in any of claims 1 to 10 and a cosmetically acceptable carrier.
12. A method for preparing the cosmetic composition according to claim 11 which comprises the step of blending lysophospholipids as defined in any of claims 1 to 10 into a cosmetic composition comprising cosmetically acceptable

carrier.

13. Use of the lysophospholipids as defined in any of claims 1 to 10 for improving the storage stability in a cosmetic composition.

5

14. Use of the lysophospholipids as defined in any of claims 1 to 10 for improving the feeling of a cosmetic composition.

15. Use of the lysophospholipids as defined in any of claims 1 to 10 for providing a suitable feeling of a cosmetic composition

10

15

20

25

30

35

40

45

50

55



European Patent  
Office

# EUROPEAN SEARCH REPORT

Application Number  
EP 99 10 5069

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (InCLs)
X	EP 0 719 541 A (INST. FOR ADVANCED SKIN RESEARCH) 3 July 1996 (1996-07-03) * page 1, line 26 - page 3, line 50; claims 1-3; examples 2,6 *	1-3,7,8, 10-12	A61K7/48 C07F9/10
X	WO 92 21323 A (PROCTER & GAMBLE) 10 December 1992 (1992-12-10) * page 4, line 20 - line 25; claims 1-4,6; example 1 *	1-3,7-12	
X	PATENT ABSTRACTS OF JAPAN vol. 12, no. 270 (C-515), 27 July 1988 (1988-07-27) & JP 63 051930 A (ASAHI DENKA), 5 March 1988 (1988-03-05) * abstract *	1	
D,Y	EP 0 255 937 A (KYOWA HAKO) 17 February 1988 (1988-02-17) * the whole document *	1-12	
Y	H. J. BURKHARDT: "Phosphatides ..." JOURNAL OF THE AMERICAN OIL CHEMISTS' SOCIETY, vol. 48, no. 11, November 1971 (1971-11), pages 697-699, XP002112745 * table III *	1-12	TECHNICAL FIELDS SEARCHED (InCLs) A61K C07F
The present search report has been drawn up for all claims			
Place of search		Date of completion of the search	Examiner
THE HAGUE		19 August 1999	Willekens, G
<p><b>CATEGORY OF CITED DOCUMENTS</b></p> <p>X : particularly relevant if taken alone  Y : particularly relevant if combined with another document of the same category  A : technological background  O : non-written disclosure  P : intermediate document</p> <p>T : theory or principle underlying the invention  E : earlier patent document, but published on, or after the filing date  O : document cited in the application  L : document cited for other reasons  A : member of the same patent family, corresponding document</p>			

EPO FORM 1503 (01.01.99) (P40001)

ANNEX TO THE EUROPEAN SEARCH REPORT  
ON EUROPEAN PATENT APPLICATION NO.

EP 99 10 5069

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

19-08-1999

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 719541	A	03-07-1996	US 5723136 A	03-03-1998
			EP 0719542 A	03-07-1996
			EP 0719543 A	03-07-1996
			WO 9535090 A	28-12-1995
			WO 9535091 A	28-12-1995
			WO 9535092 A	28-12-1995
			JP 8067619 A	12-03-1996
			JP 8067621 A	12-03-1996
			JP 8067620 A	12-03-1996
			US 5849309 A	15-12-1998
WO 9221323	A	10-12-1992	US 5238965 A	24-08-1993
			AU 2159092 A	08-01-1993
			CN 1068034 A	20-01-1993
			MX 9202557 A	01-11-1992
			PT 100562 A	30-09-1993
			TR 25911 A	01-11-1993
			US 5521223 A	28-05-1996
JP 63051930	A	05-03-1988	AT 65939 T	15-08-1991
			DE 3771923 A	12-09-1991
			EP 0245871 A	19-11-1987
			GR 3002752 T	25-01-1993
			US 4849132 A	18-07-1989
EP 255937	A	17-02-1988	JP 63041411 A	22-02-1988

EPO FORM 100/69

For more details about this annex : see Official Journal of the European Patent Office, No. 12/B2

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☒ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER: \_\_\_\_\_**

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**